


Physiological Mini Reviews

16

Volume



Vol. 16, May-June, 2023
ISSN 1669-5410 (Online)
pmr.safsiol.org.ar

Physiological
Mini
Reviews



Physiological Mini Reviews is the official journal of the Asociación Latinoamericana de Ciencias Fisiológicas, (ALACF), which is constituted by the following Societies:

Sociedad Argentina de Fisiología (SAFIS)

Sociedad Brasileira de Fisiología (SBFis)

Asociación Colombiana de Fisiología (COLFISIS)

Sociedad Cubana de Ciencias Fisiológicas (SOCCF)

Sociedad Chilena de Ciencias Fisiológicas (SCHCF)

Sociedad de Fisiología de Haití (SHF)

Sociedad Mexicana de Ciencias Fisiológicas (SMCF)

Sociedad Peruana de Ciencias Fisiológicas (SCPCF)

Sección de Fisiología de la Sociedad Uruguaya de Biociencias (SUB)

SPONSORS

We are very grateful to our sponsors

DEPARTAMENTO DE POSTGRADO
FACULTAD DE CIENCIAS MÉDICAS UNLP



maestría

EN INVESTIGACIÓN BIOMÉDICA

Acreditada y Categorizada CONEAU "C" Res. FC 2018-42
Resolución Ministerial de Reconocimiento oficial y
validez nacional del título: Res.1596/16

INSCRIPCIONES
ABIERTAS

INFORMES: maestriaibunlp@gmail.com

FACULTAD DE CIENCIAS MÉDICAS



UNIVERSIDAD NACIONAL DE LA PLATA



C I C
CONICET



LENF MEDICINA DEL SUEÑO

www.institutomedicoplatense.com.ar

Laboratorio de Electrofisiología y Medicina del Sueño.
Instituto Médico Platense. La Plata, Argentina
lenf.imp@gmail.com



intemun^{s.a}

DIVISION INSUMOS

Ventas de Insumos Médicos Hospitalarios
La Plata - Buenos Aires - Argentina
Tel: 0221 471 8951 011-75120306 0221-586 9111
ventas@intemun.com.ar administracion@intemun.com.ar

Distribuidor oficial: *Baño Fácil*



www.inbiohw.com.ar

Physiological Mini-Reviews

[ISSN 1669-5410 (Online)]

Edited by **the Argentinean Physiological Society and the Latin American Association of Physiological Sciences**

Journal address: Centro de Investigaciones Cardiovasculares y Cátedra de Fisiología y Física Biológica.
Facultad de Ciencias Médicas; Universidad Nacional de La Plata;
La Plata, Buenos Aires, Argentina. Tel.-Fax: +54-211-4834833
<http://pmr.safisiol.org.ar>

Physiological Mini-Reviews is a scientific journal, publishing brief reviews on "hot" topics in Physiology. The scope is quite broad, going from "Molecular Physiology" to "Integrated Physiological Systems". As indicated by our title it is not our intention to publish exhaustive and complete reviews. We ask to the authors concise and updated descriptions of the "state of the art" in a specific topic. Innovative and thought-provoking ideas are welcome.

Founding Editor: Mario Parisi, Buenos Aires, Argentina

Editor in Chief: Alicia Mattiazzi, La Plata, Argentina

Associate Editors

Alejandro Aiello, La Plata, Bs. As., Argentina
Walter Boron, Cleveland, OH, United States
María Jose Campagnole-Santos, Belo Horizonte, MG, Brazil
Julio Copello, Springfield, IL, United States
Ana Franchi, Ciudad Autónoma de Buenos Aires, Argentina
Cecilia Hidalgo, Santiago, Chile
Daniel Ortuño-Sahagun, Guadalajara, Jal, México
Eduardo Rios, Chicago, IL, United States
Luis Sobrevia, Santiago, Chile

Administrative Council

Presidents or delegates of Latin American Physiological Societies or Biological Societies

Ma. del Carmen Cortés Sánchez, Puebla, Pue, México
Paola Contreras, Montevideo, Uruguay
Jean-Claude Dorsainvil, Puerto Príncipe, Haití
Mayppee González Jardinez, La Habana, Cuba
Henry León, Chía, Cund., Colombia
Azael Paz Aliaga, Lima, Perú
Zully Pedroso, Santiago, Chile
Patricia Rocco, Río de Janeiro, RJ, Brazil
Martín Vila-Petroff, La Plata, Argentina

Editorial Board:

Vagner Roberto Antunes, Sao Paulo, Brazil
Cristina Arranz, Ciudad Autónoma de Buenos Aires, Argentina
Claudia Capurro, Ciudad Autónoma de Buenos Aires, Argentina
Daniel Cardinali, Ciudad Autónoma de Buenos Aires, Argentina
Marcelino Cerejido, México City, México
Alberto Crottogini, Ciudad Autónoma de Buenos Aires, Argentina
Osvaldo Delbono, Winston-Salem, NC, United States
Irene Ennis, La Plata, Bs. As., Argentina
Ariel Escobar, Merced, CA, United States
Ludmila Firaletova, St. Petersburg, Russia
Benjamín Florán Garduño, Ciudad de México, México
Ana María Gomez, Chatenay-Malabry, France
Guillermo González Burgos Pittsburg, PA, United States
Carlos González, Lubbock, TX, United States
Hilda Leonor González Olaya, Bucaramanga, SAN, Colombia

Sergio Gradilone, Rochester, MN, United States
Joost Hoenderop, Nijmegen, The Netherlands
Bredford Kerr Fuentes, Santiago, Chile.
Cecilia Larocca, Rosario, Santa Fe, Argentina
Elena Lascano, Ciudad Autónoma de Buenos Aires, Argentina
Reinaldo Marín, Caracas, Venezuela
Raúl Marinelli, Rosario, Santa Fé, Argentina
Susana Mosca, La Plata, Bs. As., Argentina
Cecilia Mundiña-Weilemann, La Plata, Bs. As., Argentina
Gustavo Pérez, La Plata, Bs. As., Argentina
Darío Protti, Sidney, NSW, Australia
Margarita Salas, La Plata, Bs. As., Argentina
Daniel Schulz, Gif sur Yvette, France
Gary Sieck, Rochester, MN, United States

Education Editorial Board

Robert G. Carroll, Greenville, NC, United States
Matilde Said, La Plata, Bs. As., Argentina
Roxana Troiano, Ciudad Autónoma de Buenos Aires, Argentina
Claudia Caldiz, La Plata, Bs. As., Argentina

Publishing Editorial Board

Carolina Caniffi, Ciudad Autónoma de Buenos Aires, Argentina
Verónica de Giusti, La Plata, Bs. As., Argentina
Luis Gonano, La Plata, Bs. As., Argentina
Alejandro Orlowsky, La Plata, Bs. As., Argentina

Zully Pedroso, Santiago, Chile
Matilde Said, La Plata, Bs. As., Argentina
Carlos Valverde, La Plata, Bs. As., Argentina
Alejandra Yeves, La Plata, Bs. As., Argentina

Editorial Assistant: María Ines Vera

Preparation and Submission of manuscripts:

"Physiological Mini-Reviews" will have a maximum of 3000 words, 50 references and 3 figures. Material will be addressed to scientific people in general but not restricted to specialist of the field. For citations in the text please refer to Instructions in our webpage. Final format will be given at the Editorial Office. Most contributions will be invited ones, but spontaneous presentations are welcome. Send your manuscript in Word format (.doc or .docx) to: pmr.alacf@gmail.com

Advertising: For details, rates and specifications contact the Associate Editor at the Journal address e-mail: pmr.alacf@gmail.com

INNOVATIVE TECHNIQUES AND NEW INSIGHTS

Studying cardiac ionic currents and action potentials in physiologically relevant conditions

Ye Chen-Izu ^{1,2,3}, Bence Hegyi ¹, Zhong Jian ¹, Balazs Horvath ^{1,4}, John A. Shaw ^{1,5},
Tamas Banyasz ^{1,4}, Leighton T. Izu ^{1*}

¹ Department of Pharmacology, University of California, Davis, USA

² Department of Biomedical Engineering, University of California, Davis, USA

³ Department of Internal Medicine, Division of Cardiovascular Medicine, University of California, Davis, USA

⁴ Department of Physiology, University of Debrecen, Hungary

⁵ Department of Aerospace Engineering, University of Michigan, Ann Arbor, USA

*correspondence to: LTizu@ucdavis.edu

Abstract

Cardiac arrhythmias are associated with various forms of heart diseases. Ventricular arrhythmias present a significant risk for sudden cardiac death. Atrial fibrillations predispose to blood clots leading to stroke and heart attack. Scientists have been developing patch-clamp technology to study ion channels and action potentials (APs) underlying cardiac excitation and arrhythmias. Beyond the traditional patch-clamp techniques, innovative new techniques were developed for studying complex arrhythmia mechanisms. Here we review the recent development of methods including AP-Clamp, Dynamic Clamp, AP-Clamp Sequential Dissection, and Patch-Clamp-in-Gel. These methods provide powerful tools for researchers to decipher how the dynamic systems in excitation-Ca²⁺ signaling-contraction feedforward and feedback to control cardiac function and how their dysregulations lead to heart diseases.

Keywords: action potential, arrhythmia, cardiac myocyte, patch-clamp, electrophysiology.

Resumen

Las arritmias cardiacas están asociadas a diferentes tipos de enfermedad cardiaca. Las arritmias ventriculares constituyen un alto riesgo de muerte súbita. La fibrilación auricular predispone a coágulos sanguíneos que pueden producir accidentes cerebrovasculares e infarto miocárdico. Los científicos han desarrollado la técnica de patch-clamp para estudiar los canales iónicos y los potenciales de acción (PAs), que constituyen la base de la excitación y las arritmias cardiacas. Además de las clásicas técnicas de patch-clamp, se desarrollaron técnicas innovativas para estudiar los mecanismos complejos de las arritmias. En este trabajo, describimos diferentes métodos recientemente desarrollados tales como AP-clamp (“clampeo” del PA), Dynamic Clamp (“clampeo” dinámico), AP-Clamp Sequential Dissection, (disección secuencial del “clampeo” del AP), y Patch-Clamp-in-Gel (Patch clamp en gel). Estos métodos constituyen herramientas poderosas para descifrar cómo los sistemas dinámicos que constituyen la excitación-las señales de Ca²⁺ y la contracción, se retroalimentan para controlar la función cardiaca y cómo sus alteraciones llevan a la enfermedad cardiaca.

Palabras clave: potencial de acción, arritmias, miocitos cardiacos, patch -clamp, electrofisiología.

Introduction

Cardiac action potentials drive the rhythmic heartbeats and dysregulation of action potentials can lead to cardiac arrhythmias. In cardiomyocytes, the action potential (AP) is determined by myriad ionic currents that work in concert to control the rise and fall of the transmembrane voltage. The AP timing and voltage profile is dynamically shaped by the counterbalancing effects of inward currents versus outward currents. Researchers have developed various patch-clamp techniques to investigate ion channels and action potentials to understand their roles in cell function and diseases. Traditionally, the voltage-clamp technique is used to determine ion channel properties and the current-clamp technique is used to record the cell's action potential. In this minireview, we will focus on the recent development of innovative techniques for studying the dynamic ionic currents under the AP in physiologically relevant conditions that closely mimic the in-situ environment for studying how the dynamic ionic currents contribute to shaping the action potential and arrhythmogenic activities.

AP-Clamp techniques to study the dynamics of ionic currents during cardiac AP cycle

To study the dynamics of ionic currents during the AP cycle, AP-clamp techniques were developed by using the AP wave form to voltage-clamp the cell. [5, 6, 7, 8, 9] In comparison, traditional voltage-clamp typically uses a rectangular voltage pulse to record the current flowing through a particular ion channel and study its electrophysiological properties (i.e., ion selectivity, conductance, gating, activation, inactivation, recovery, etc.) at constant voltage. These properties were then put into a mathematical model using the approach pioneered by Hodgkin–Huxley [10], which is then used to calculate the profile of the ionic currents during an AP. The AP-clamp technique directly measures the ionic current during the AP using the following experimental protocol (Figure-1A): (1) Under current-clamp ($I = 0$) mode, pace the cell to reach steady-state and record the cell's steady-state AP. (2) Switch to voltage-clamp mode, and use this AP waveform as the voltage command to voltage-clamp the cell and repeat the cycle until reaching steady-state; the net current output – the sum of all currents - should be zero at steady-state, which serves as the reference current (I_{Ref}). (3) Block an ionic current by using its specific blocker to remove it from the net current output; this missing current is seen in the compensation current (I_{Comp}) injected by the amplifier to maintain the AP waveform. (4) The difference between the reference and the compensation current ($I_{Diff} = I_{Ref} - I_{Comp}$) shows the ionic current that was originally flowing during the AP prior to the blocker application. The above protocol is used with some variations: canonical AP-Clamp using a pre-recorded AP waveform to clamp different cells [11], reconstructed AP-Clamp using a reconstructed AP waveform to study the ion channel kinetic properties [12], or self AP-Clamp using the AP recorded from the same cell to do AP-clamp to mimic the physiological condition in the cell. [3, 13].

There are fundamental differences between the AP-clamp techniques and the traditional voltage-clamp techniques. The AP-clamp records an ionic current over a changing voltage during an AP, whereas the traditional voltage-clamp records the current at constant voltage. AP-clamp experiments record the 'absence of the specific current' by blocking one single current using its specific blocker while allowing all other currents to flow during AP in physiological condition. In contrast, traditional voltage-clamp experiments record a single current using non-physiological condition that suppress all other currents (i.e., using rectangular pulse voltage protocols, simplifying solutions to select ion species, blocking other contaminating currents using pharmacological inhibitors, etc.) Furthermore, the intracellular Ca^{2+} transient during AP-clamp can be preserved by eliminating exogenous Ca^{2+} buffer in the pipette solution to mimic the physiological condition in cardiomyocytes.

AP-Clamp allows direct recording of the ionic currents during the dynamically changing action potential. For example, Horvath et al. [14] recorded the TTX-sensitive Na^+ current under AP-Clamp, which shows a distinct profile that clearly separates the late Na^+ current (I_{NaL}) from the fast Na^+ current. The I_{NaL} profile under AP-Clamp drastically differs from the I_{NaL} recorded under rectangular pulse voltage-clamp where the I_{NaL} merges with the fast Na^+ current and the two are hard to separate [15]. The AP-Clamp data reveals that the I_{NaL} during cardiac AP is much larger than previously estimated. Using AP-Clamp, Banyazs et al. [16] discovered that the L-type Ca^{2+} channel (I_{Ca-L}) has

rapid deactivation and reactivation during the AP notch; this causes transmural difference in I_{Ca-L} because the epicardium has a more pronounced AP notch than the endocardium. Combined with Ca^{2+} measurement, AP-clamp with Ca^{2+} cycling provides a powerful method for studying the dynamic interplay of Ca^{2+} , ionic currents, action potentials and arrhythmogenic activities. [11, 13, 17]. For example, Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII) is an important regulator of ion channels including I_{Ca-L} and I_{NaL} . AP-clamp with Ca^{2+} cycling experiments revealed the dynamic profile of I_{NaL} during cardiac AP and its prominent role in arrhythmogenesis. [14, 18] The AP-clamp with Ca^{2+} cycling method also enables studying how intracellular Ca^{2+} transient modulates the Ca^{2+} -sensitive currents, such as the L-type Ca^{2+} current and the Na / Ca^{2+} exchanger current, during the cardiac AP cycle. [13]




Figure 1. AP-Clamp method for studying the dynamics of ionic currents during cardiac AP.

(Figures reconstructed from Banyasz et al., 2011 [3].)

- A. Schematic of AP-clamp technique: (1) Record the cell's steady state AP. (2) Use this AP as the voltage clamp command to record the total net current output, I_{Ref} . (3) Isolate a current by using its specific blocker to remove it from the total net current output, and record the compensation current from the amplifier, I_{Comp} . (4) The current of interest is obtained as the difference current: $I_{Diff} = I_{Ref} - I_{Comp}$.
- B. The first current I_{Ks} is obtained by subtracting the compensation current recorded with chromanol-293B from the baseline current.
- C. The second current I_{Kr} is obtained by subtracting the compensation current recorded with E4031+chromanol from the first compensation current recorded with chromanol.
- D. AP-Clamp Sequential Dissection recording of 4 currents, I_{Ks} , I_{Kr} , I_{K1} and I_{Ca-L} using the corresponding channel blockers chromanol-293B, E4031, Ba^{2+} and nisoldipine.
- E. AP-Clamp Sequential Dissection recording of 4 currents, I_{Ks} , I_{Kr} , I_{K1} and I_{Ca-L} from another cell, showing smaller current densities in all currents.
- F. Correlation between the charges carried by inward I_{Ca-L} and outward I_{Ks} .
- G. Correlation between the charges carried by inward I_{Ca-L} and the sum of charges carried by the outward I_{Ks} , I_{Kr} , I_{K1} currents.

Dynamic Clamp for studying dynamic interaction of the AP and the underlying currents

The Dynamic Clamp method was developed to investigate the interplay between ionic currents and action potentials in cardiomyocytes. The Dynamic Clamp electrically connects two cells (cell-A, cell-B) to allow intercellular electrical coupling. The ionic current in cell-B can be manipulated genetically or pharmacologically, or custom designed using computational model simulation. This current from cell-B is injected into cell-A to study its effect on modifying the cell's action potential. In the recent decades, Dynamic Clamp has evolved to include diverse techniques such as coupling clamp, model clamp, and dynamic AP-Clamp. [19] The technical implementations of Dynamic Clamp require sophisticated hardware and software, which had been nicely summarized in previous reviews. [19, 20] Here we briefly mention the advantages as well as limitations of the method.

Dynamic Clamp provides powerful tools to manipulate the injected current to study its effect on modifying the action potential in real-time. The injected current can be recorded from another cell expressing a native or mutant ion channel or from a mathematical model simulation. For example, Berecki et al. [21] used coupling clamp to study how various mutations in the rapid rectifier K^+ current (I_{Kr}) affect the AP. They removed native I_{Kr} using specific pharmacological blocker E-4031 in a cardiomyocyte (cell-A). They expressed mutant I_{Kr} in a HEK-293 cell (cell-B), recorded the mutant I_{Kr} current under AP-clamp (using the AP from cell-A), and injected it into cell-A. This intercellular coupling incorporates the mutant I_{Kr} into the AP in real time. They found that wild-type I_{Kr} injection restored the original APD, but R56Q mutation prolonged APD leading to long-QT2 syndrome. Using this approach, they also found that cardiac Na^+ channel (SCN5A) mutations Y1795C and A1330P prolonged APD leading to long-QT3 syndrome. [22] Using the model clamp approach, Altomare et al. [23] blocked the native I_{Kr} in a cardiomyocyte (cell-A); generated an artificial current using a computational model of I_{Kr} (cell-B); and then injected the artificial current to study its effect on the AP profile. They found that the model generated I_{Kr} current could adequately replace the native I_{Kr} to recover the original APD, attesting to the accuracy of the I_{Kr} model. Then they manipulated the model parameters to study the effect of each parameter on modifying the I_{Kr} and the AP profile. They found that the APD variability is sensitive to the steady-state inactivation more than to the activation. This approach enables systematic and detailed studies on how the voltage-time dependent channel properties influence the current and the AP. Hence, Dynamic Clamp providing a powerful tool to simulate the effects of mutations or drugs on modifying the channel properties and the consequences on the AP and arrhythmogenic activities.

Dynamic Clamp also allows investigating the mechanistic details in the interplay between AP profile and ionic currents. Experiments injecting I_{to} to manipulate AP notch revealed that the 'spike-and-dome' and the plateau height critically determine the AP duration, which demystifies why different animal species with different AP profiles have differential response to β -adrenergic stimulation. [24] Dynamic Clamp experiments also showed that the β -adrenergic upregulation of the funny current (I_f) in cardiac pacemaker cells accounts for 60% of the AP interval decrease and 41% of the AP firing rate increase for the fight-or-flight response. [25] Recently, Dynamic Clamp is used to inject an artificial I_{K1} current into human induced pluripotent stem cell-derived cardiomyocytes (hiPSC-CMs, with immature AP profile) to simulate "adult-like" AP response to drugs. [26]

A major limitation of the Dynamic Clamp method is due to that the injected current uses electron as the charge carrier, which differs from the ion species (i.e., Na^+ , K^+ , Ca^{2+} , etc.) that flow through selective ion channels. The ionic currents are necessary for generating the ionic gradients that set the electrochemical potentials across the cell membrane. Restoring the ionic homeostasis is needed for maintaining the cell function during cardiac cycle. However, the roles of ionic currents in establishing ionic homeostasis cannot be mimicked by the electron injection used in Dynamic Clamp.

AP-Clamp Sequential Dissection to study the counterbalance of ionic currents during AP

AP-clamp Sequential Dissection technique, nicknamed 'Onion-Peeling', was further developed from AP-Clamp to record many different ionic currents in the same cell. [3, 1] Previous patch-clamp techniques record only one current in any one cell. The AP-clamp Sequential Dissection uses a series

of channel blockers to sequentially dissect out many ionic currents one by one in a single cell under self AP-Clamp. The experimental protocol mainly consists of the following steps. First, the cell is placed under AP-clamp; a specific blocker is applied to dissect out its corresponding ionic current (Figure-1B). Next, a second blocker is applied to dissect out the second current (Figure-1C). Next, a third blocker is applied to dissect out the third current, etc. Figure-1D shows 4 ionic currents (I_{Ks} , I_{Kr} , I_{K1} , I_{Ca-L}) dissected out by sequentially applying the specific blockers for each current. [3] A technical limitation for the AP-Clamp Sequential Dissection method is the specificity of channel blockers. To minimize the off targets effects, one can use low dosage of blockers that is usually more specific, and then scale to the full current density. To preserve the Ca^{2+} transient during ‘onion-peeling’, the Ca^{2+} channel blocker is used only at the end after other channel blockers.

The unprecedented ability to measure many different currents in the same cell enables studying how the inward currents versus the outward currents counterbalance to shape the cell’s AP. Studies using this technique show that the contribution of different ionic currents to shaping the AP may dramatically change under various conditions such as beta-adrenergic stimulation or heart disease development. [27, 1] AP-clamp Sequential Dissection can be used to study many different ionic currents and integrate them at the individual cell level, which provides a comprehensive view on the individual cell electrophysiology as well cell-to-cell variability at the tissue level.

Patch-Clamp-in-Gel to study the mechanical load regulation of AP and ionic currents

The heart pumps blood against the mechanical load from arterial resistance, and increased load may alter cardiac electrophysiology and contribute to arrhythmias. To study mechanical load effects on cardiomyocytes, the Cell-in-Gel technology was developed to embed isolated cardiomyocytes in a 3-D hydrogel made of viscoelastic polymer matrix. [28] Mechanical analyses show that when the cell is contracting in-gel, the viscoelastic gel exerts 3-D mechanical stress on the cell (Figure-3A, longitudinal tension, transverse compression, and surface traction with normal and shear stress). [2, 29, 30, 31] To study the mechanical load effects on electrophysiology, the Patch-Clamp-in-Gel technique was further developed to control the mechanical load on the cell in patch-clamp experiments. [4] Figure-3B shows a schematic of the Patch-Clamp-in-Gel technique. First, perform standard patch-clamp experiment (whole-cell configuration) in a cell bathed in Tyrode’s solution containing polyvinyl alcohol (PVA). Pace the cell to reach steady-state and record the APs (Figure-3C, load-free control). Next, add tetraboronate-polyethylene glycol (4B-PEG) to crosslink PVA to form polymer matrix that embeds the cell-electrode assembly in a viscoelastic hydrogel (Figure-3D, cell in-gel). As the hydrogel forms gradually over minutes, the AP duration (APD_{90}) initially became longer in the cell contracting under moderate load in softer gel (Figure-3E, 3G), followed by development of APD alternans in the cell contracting under higher load in stiffer gel (Figure-3F, 3G). These data demonstrate that the mechanical load on cardiomyocytes during cell contraction profoundly regulates the cell’s electrophysiological properties.

Patch-Clamp-in-Gel experiments show that the cardiomyocyte under mechanical load has increased L-type Ca^{2+} current, increased transient outward K^+ current, decreased inward rectifier K^+ current, and prolonged APD. [4] These changes allow more Ca^{2+} to enter the cell during the AP cycle, causing elevated intracellular Ca^{2+} content that increases the SR Ca^{2+} release and cytosolic Ca^{2+} transient. Increased Ca^{2+} transient gives rise to enhanced contractility of the cardiomyocyte contracting under mechanical load. [31, 32] Taken together, these experimental data reveal the mechano-chemo-electro-transduction (MCET) mechanisms that enable cardiomyocytes to upregulate contractility to compensate for the increased load. Furthermore, systematic studies applying different levels of mechanical load on cardiomyocytes show that MCET provides feedback loops in the dynamic system of excitation- Ca^{2+} signaling-contraction (E-C) coupling, which enables autoregulation of E-C coupling and contractility in cardiomyocytes. [31, 32] Studies also show that exceedingly high load causes dysregulation of MCET, spontaneous Ca^{2+} activities, and discordant alternans in APD and Ca^{2+} transient, which provide substrate to cardiac arrhythmias. [4, 28, 32]

New insights on the complex changes of cellular electrophysiology in health and diseases

Cardiac action potential is shaped by myriad ionic currents flowing in or out of the cell in a time-dependent manner. The AP-clamp Sequential Dissection method enables recording many different currents from a single cell, which is a significant step beyond the traditional way of recording only one current in any one cell. The ability to measure different currents in a single cell revealed hitherto unknown characteristics of the ionic currents in individual cardiomyocytes: coordination of the currents within a cell and large variations between different cells. In the first Onion-Peeling study [3], four different ionic currents (I_{Ks} , I_{Kr} , I_{K1} , and I_{Ca-L}) were measured from [21] ventricular myocytes isolated from healthy guinea pig heart. The data reveal that in a single cell, the inward currents and the outward currents are coordinated to produce a normal AP profile. However, different cells have large variations in the amplitudes of currents (Figure-1D, 1C). Interestingly, despite the large cell-to-cell variation (Figure-1F, 1G), the changes in the inward and the outward currents in each cell are coordinated to keep the normal AP profile across the individual cells in the healthy heart.

In failing hearts, the coordination between ionic currents may breakdown. Study of the ischemic heart failure (HF) [1] shows that the ionic currents in cardiomyocytes from the infarct border and remote zones undergo complex remodeling in HF (Figure-2A). The ventricular myocytes isolated from different zones show different action potentials (Figure-2B). AP-Clamp Sequential Dissection was used to record eight different ionic currents during the cell's AP under physiologically relevant condition (Figure-2C). Compared with healthy control, infarct-remote zone myocytes showed increases in the late Na^+ current, the Ca^{2+} -activated K^+ current, the Ca^{2+} -activated Cl^- current, but a decrease of the rapid delayed rectifier K^+ current and an altered Na^+/Ca^{2+} exchange current profile (Figure-2D, 2E). In the infarct-border zone myocytes, additional changes occurred with a decrease of the L-type Ca^{2+} current and a decrease of the inward rectifier K^+ current (Figure-2D, 2F). The changes in any individual current are relatively small, but the integrated impacts shift the balance between the inward currents and the outward currents to shorten AP in the border zone but prolong AP in the remote zone. Such differential remodeling increases the heterogeneity and dispersion of action potential durations, providing substrates to cardiac arrhythmias. Another study using AP-clamp Sequential Dissection shows that I_{Kr} and I_{NaL} are counterbalancing currents during the AP and they covary in a concerted manner in individual ventricular myocytes. [33] Moreover, the ratio of I_{Kr}/I_{NaL} is altered in HF, long QT and short QT syndromes, and restoring the I_{Kr}/I_{NaL} balance may represent a therapeutic strategy. Hence, the AP-clamp Sequential Dissection method provides a unique and powerful tool for studying the complex changes of cardiac electrophysiology in heart diseases as well as the complex effects of drug therapies.

In conclusion, the innovative techniques reviewed here provide powerful tools for investigating the complex mechanisms of cardiac arrhythmias. The AP-clamp techniques allow direct recording of the dynamic ionic currents during the action potential in physiologically relevant conditions. Dynamic Clamp provides versatile tools to manipulate ionic currents to study how they affect the AP profile. AP-Clamp Sequential Dissection enables recording of many different ionic currents in the same cell to study how they integrate to shape the cell's AP. Patch-Clamp-in-Gel provides a new tool for studying how the mechanical load during cell contraction feedback to autoregulate the E-C coupling systems. Studying E-C coupling as an autoregulatory feedback system will help deciphering how the heart autoregulates contractility in response to physiological load changes, and how pathological overload such as hypertension leads to cardiac arrhythmias and heart failure.




Figure 2. AP-Clamp Sequential Dissection to study complex changes in electrophysiology.

(Figures reconstructed from Hegyi et al., 2018 PNAS¹.)

(A) Ischemic heart failure (HF) showing the infarct, border zone, and remote zone.

(B) Representative APs from control, HF-remote, and HF-border cardiomyocytes at 36°C.

(C) AP-Clamp Sequential Dissection recording of 8 ionic currents from the same cell.

Dynamic ionic currents during the AP were recorded from the control (D), the HFremote zone, and the HF-border zone (F). The contribution of each inward and outward current was normalized to the momentary total outward (RD) or inward (DD) membrane current calculated based on the measured eight ionic current densities. Ionic currents measured in each cell were also normalized to the corresponding APD₉₅. Bottom panels show the fingerprinting of ionic currents using AP-Clamp Sequential Dissection.




Figure 3. Patch-Clamp-in-Gel to study mechanical load effect on regulating ion channels and AP. (Figures reconstructed from Kazemi-Lari et al., 2021[2]; and Hegyi et al., 2021[4].) (A) Schematic of the Patch-Clamp-in-Gel method. (B) Cell-in-Gel system applies 3-D mechanical stress on the cardiomyocyte during contraction. (C) Patch-clamp experiments were performed in a cardiomyocyte in Tyrode solution containing PVA (load-free control). (D) Then, 4B-PEG cross-linker was added to form a 3-D hydrogel around the cell to apply mechanical load during cell contraction. (E) APD was prolonged in the cell contracting under moderate load in softer gel. (F) APD alternans developed in the cell contracting under higher load in stiffer gel. (G) APD₉₀ changes during the formation of hydrogel that progressively increase the mechanical load on the cell.

References

- [1] **Hegyi B, Bossuyt J, Griffiths LG, Shimkunas R, Coulibaly Z, Jian Z, Grimsrud KN, Sondergaard CS, Ginsburg KS, Chiamvimonvat N et al.** Complex electrophysiological remodeling in postinfarction ischemic heart failure. *Proc Natl Acad Sci U S A.* 2018;115(13):E3036-E44. Epub 2018/03/14.
- [2] **Kazemi-Lari MA, Shimkunas R, Jian Z, Hegyi B, Izu L, Shaw JA, Wineman AS, Chen-Izu Y.** Modeling cardiomyocyte mechanics and autoregulation of contractility by mechano-chemo-transduction feedback. *iScience.* 2022;25(7):104667. Epub 20220626.
- [3] **Banyasz T, Horvath B, Jian Z, Izu LT, Chen-Izu Y.** Sequential dissection of multiple ionic currents in single cardiac myocytes under action potential-clamp. *J Mol Cell Cardiol.* 2011;50(3):578-81. Epub 2011/01/11.
- [4] **Hegyi B, Shimkunas R, Jian Z, Izu LT, Bers DM, Chen-Izu Y.** Mechanoelectric coupling and arrhythmogenesis in cardiomyocytes contracting under mechanical afterload in a 3D viscoelastic hydrogel. *Proc Natl Acad Sci U S A.* 2021;118(31). Epub 2021/07/31.
- [5] **Starzak ME, Starzak RJ.** An action potential clamp to probe the effectiveness of space clamp in axons. *IEEE Trans Biomed Eng.* 1978;25(2):201-4.
- [6] **Doerr T, Denger R, Trautwein W.** Calcium currents in single SA nodal cells of the rabbit heart studied with action potential clamp. *Pflugers Arch.* 1989;413(6):599-603.
- [7] **de Haas V, Vogel W.** Sodium and potassium currents recorded during an action potential. *European biophysics journal: EBJ.* 1989;17(1):49-51.
- [8] **Bouchard RA, Clark RB, Giles WR.** Effects of action potential duration on excitation-contraction coupling in rat ventricular myocytes. Action potential voltage-clamp measurements. *Circ Res.* 1995;76(5):790-801. Epub 1995/05/01.
- [9] **Grantham CJ, Cannell MB.** Ca²⁺ Influx During the Cardiac Action Potential in Guinea Pig Ventricular Myocytes. *Circulation Research.* 1996;79(2):194.
- [10] **Hodgkin ALaH, A. F.** A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol.* 1952;117:500-44.
- [11] **Arreola J, Dirksen RT, Shieh RC, Williford DJ, Sheu SS.** Ca²⁺ current and Ca²⁺ transients under action potential clamp in guinea pig ventricular myocytes. *Am J Physiol.* 1991;261(2 Pt 1):C393-7.
- [12] **Varro A, Balati B, Iost N, Takacs J, Virag L, Lathrop DA, Csaba L, Talosi L, Papp JG.** The role of the delayed rectifier component I_{Ks} in dog ventricular muscle and Purkinje fibre repolarization. *J Physiol.* 2000;523 Pt 1(Pt 1):67-81.
- [13] **Banyasz T, Horvath B, Jian Z, Izu LT, Chen-Izu Y.** Profile of L-type Ca²⁺ current and Na⁺/Ca²⁺ exchange current during cardiac action potential in ventricular myocytes. *Heart Rhythm.* 2012;9(1):134-42.
- [14] **Horvath B, Banyasz T, Jian Z, Hegyi B, Kistamas K, Nanasi PP, Izu LT, Chen-Izu Y.** Dynamics of the late Na⁺ current during cardiac action potential and its contribution to afterdepolarizations. *Journal of Molecular and Cellular Cardiology.* 2013;64(0):59-68.
- [15] **Rajamani S, El-Bizri N, Shryock JC, Makielski JC, Belardinelli L.** Use-dependent block of cardiac late Na(+) current by ranolazine. *Heart Rhythm.* 2009;6(11):1625-31.

- [16] **Banyasz T, Magyar J, Szentandrassy N, Horvath B, Birinyi P, Szentmiklosi J, Nanasi PP.** Action potential clamp fingerprints of K⁺ currents in canine cardiomyocytes: their role in ventricular repolarization. *Acta Physiol (Oxf)*. 2007;190(3):189-98.
- [17] **Lyu Y, Thai PN, Ren L, Timofeyev V, Jian Z, Park S, Ginsburg KS, Overton J, Bossuyt J, Bers DM et al.** Beat-to-beat dynamic regulation of intracellular pH in cardiomyocytes. *iScience*. 2022;25(1):103624. Epub 20211213.
- [18] **Hegy B, Morotti S, Liu C, Ginsburg KS, Bossuyt J, Belardinelli L, Izu LT, Chen-Izu Y, Banyasz T, Grandi E et al.** Enhanced Depolarization Drive in Failing Rabbit Ventricular Myocytes. *Circ Arrhythm Electrophysiol*. 2019;12(3):e007061. Epub 2019/03/19.
- [19] **Wilders R.** Dynamic clamp: a powerful tool in cardiac electrophysiology. *J Physiol*. 2006;576(Pt 2):349-59. Epub 20060727.
- [20] **Ortega FA, Grandi E, Krogh-Madsen T, Christini DJ.** Applications of Dynamic Clamp to Cardiac Arrhythmia Research: Role in Drug Target Discovery and Safety Pharmacology Testing. *Frontiers in physiology*. 2017;8:1099. Epub 20180104.
- [21] **Berecki G, Zegers JG, Verkerk AO, Bhuiyan ZA, de Jonge B, Veldkamp MW, Wilders R, van Ginneken AC.** HERG channel (dys)function revealed by dynamic action potential clamp technique. *Biophys J*. 2005;88(1):566-78. Epub 20041008.
- [22] **Berecki G, Zegers JG, Bhuiyan ZA, Verkerk AO, Wilders R, van Ginneken AC.** Long-QT syndrome-related sodium channel mutations probed by the dynamic action potential clamp technique. *J Physiol*. 2006;570(Pt 2):237-50. Epub 20051027.
- [23] **Altomare C, Bartolucci C, Sala L, Bernardi J, Mostacciolo G, Rocchetti M, Severi S, Zaza A.** IKr Impact on Repolarization and Its Variability Assessed by Dynamic Clamp. *Circ Arrhythm Electrophysiol*. 2015;8(5):1265-75.
- [24] **Sala L, Hegyi B, Bartolucci C, Altomare C, Rocchetti M, Vaczi K, Mostacciolo G, Szentandrassy N, Severi S, Pal Nanasi P et al.** Action potential contour contributes to species differences in repolarization response to beta-adrenergic stimulation. *Europace : European pacing, arrhythmias, and cardiac electrophysiology : journal of the working groups on cardiac pacing, arrhythmias, and cardiac cellular electrophysiology of the European Society of Cardiology*. 2018;20(9):1543-52.
- [25] **Peters CH, Rickert C, Morotti S, Grandi E, Aronow KA, Beam KG, Proenza C.** The funny current I_f is essential for the fight-or-flight response in cardiac pacemaker cells. *J Gen Physiol*. 2022;154(12). Epub 20221028.
- [26] **Verkerk AO, Wilders R.** Dynamic Clamp in Electrophysiological Studies on Stem Cell-Derived Cardiomyocytes-Why and How? *J Cardiovasc Pharmacol*. 2021;77(3):267-79.
- [27] **Banyasz T, Jian Z, Horvath B, Khabbaz S, Izu L, Chen-Izu Y.** Beta-adrenergic stimulation reverses the I_{Kr}-I_{Ks} dominant pattern during cardiac action potential. *Pflugers Arch - Eur J Physiol*. 2014;466(11):2067-76.
- [28] **Jian Z, Han H, Zhang T, Puglisi J, Izu LT, Shaw JA, Onofriok E, Erickson JR, Chen Y-J, Horvath B et al.** Mechanochemotransduction During Cardiomyocyte Contraction Is Mediated by Localized Nitric Oxide Signaling. *Sci Signal*. 2014;7(317):ra27-.
- [29] **Shaw J, Izu L, Chen-Izu Y.** Mechanical Analysis of Single Myocyte Contraction in a 3-D Elastic Matrix. *PLoS ONE*. 2013;8(10):e75492.

- [30] **Kazemi-Lari MA, Shaw JA, Wineman AS, Shimkunas R, Jian Z, Hegyi B, Izu L, Chen-Izu Y.** A viscoelastic Eshelby inclusion model and analysis of the Cell-in-Gel system. *Int J Eng Sci.* 2021;165. Epub 2021/10/12.
- [31] **Izu L, Shimkunas R, Jian Z, Hegyi B, Kazemi-Lari M, Baker A, Shaw J, Banyasz T, Chen-Izu Y.** Emergence of Mechano-Sensitive Contraction Autoregulation in Cardiomyocytes. *Life (Basel).* 2021;11(6). Epub 2021/06/03.
- [32] **Shimkunas R, Hegyi B, Jian Z, Shaw JA, Kazemi-Lari MA, Mitra D, Leach JK, Li X, Jaradeh M, Balardi N et al.** Mechanical Load Regulates Excitation-Ca(2+) Signaling-Contraction in Cardiomyocyte. *Circ Res.* 2021;128(6):772-4. Epub 2021/02/20.
- [33] **Hegyi B, Chen-Izu Y, Izu LT, Rajamani S, Belardinelli L, Bers DM, Banyasz T.** Balance Between Rapid Delayed Rectifier K(+) Current and Late Na(+) Current on Ventricular Repolarization: An Effective Antiarrhythmic Target? *Circ Arrhythm Electrophysiol.* 2020;13(4):e008130. Epub 2020/03/24.


ABOUT AUTHORS



Dr. Ye Chen-Izu, MS, PhD, FAHA, is a Professor of Bioengineering, Pharmacology, and Cardiology at the University of California, Davis, USA. As the Director of the Cardiac Signaling Lab, she leads an interdisciplinary team to investigate heart disease mechanisms and develop new therapeutic strategies. She is an elected Fellow of the American Heart Association (FAHA) and served as a Regular Member in the National Institutes of Health (NIH) grants review sections. She has received numerous grants and awards and published 80+ scientific research articles, book chapters, and teaching materials. Dr. Chen-Izu is a well-recognized innovator who has developed a series of new technologies to study how mechanical load such as high blood pressure affects cardiac excitation-contraction coupling and heart diseases. She also leads translational study to develop High-content High-throughput (HCHT) Cardiotoxicity Screening technology pharmaceutical drug R&D.



Tamas Banyasz is a professor at Department of Physiology, Medical and Health Science Center, University of Debrecen, Debrecen, Hungary. Received a medical degree from the University Medical School of Debrecen in 1984. He has PhD on Physiology and Neurobiology (1996) and D.Sc. from Hungarian Academy of Science (2017). He spent several years as a visiting scientist at various research labs (University of Oklahoma; Universita Degli Studi Milano Bicocca; University of Kentucky, University of California Davis). He publishes on the field of cardiac physiology and pharmacology with a focus on cellular electrophysiology, cellular calcium homeostasis and cardiac arrhythmogenesis. He has been involved in multiple international collaborative research projects in the field. He authors 132 research papers and 4 book chapters related to his research activity that resulted in more than 2000 independent citations. He is Editorial Board member at *Frontiers in Physiology* (EISSN 1664-042X) and served as reviewer on more than 20 journals. He is lecturer of Physiology at Faculties of General Medicine, Dentistry and Pharmacology at University of Debrecen and program coordinator of several courses. He co-authored 15 university textbooks.



Zhong Jian, Ph.D. with a multidisciplinary education and training background in semiconductor physics (B.S), statistical physics (M.S), and biophysics (Ph.D.), Zhong Jian has established a strong foundation in interdisciplinary research and expertise, particularly in the realm of cardiovascular disease mechanisms.

His research has been devoted to investigating the cellular signaling mechanisms underlying cellular mechanosensitivity and the resulting pathological consequences. His studies have encompassed various areas, such as exploration of neuronal dynamics in chronic compression-induced pathological pain, examination of TRPC channel gating properties and their mechanosensitivity, investigation of the ionic mechanisms of arrhythmia, and exploration of Ca²⁺ signaling in cardiac excitation-contraction coupling relevant to heart disease mechanisms. In particular, over the past decade, he has focused specifically on studying mechano-chemo-electro transduction (MCET), which aims to understand the feedback from contraction to the Ca²⁺ signaling system and electrical system in cardiomyocytes. To achieve this, he has developed innovative technologies such as the patch-clamp-in-gel technique and in vivo cannulation methods for isolating cardiomyocytes from heart disease models.



Bence Hegyi, MD, PhD, is currently a research scientist at the Department of Pharmacology, University of California, Davis. His research focuses on studying various molecular and cellular mechanisms responsible for the development of cardiac arrhythmias and contractile dysfunctions in heart failure and diabetes with a clear scope of linking molecular alterations to disease and translating basic research findings to clinics. He received a rigorous training in state-of-the-art electrophysiology and fluorescent imaging techniques (under the mentorship of Drs. Ye Chen-Izu and Donald M. Bers during his postdoc period), and successfully combined these -individually challenging- techniques to better understand the feedback interactions between the electrical, the Ca²⁺ signaling, and the contractile systems in cardiomyocytes under physiological conditions. He also significantly contributed to establish and develop several animal models of diabetic cardiomyopathy and heart failure, including new translational mouse models for heart failure with preserved ejection fraction (HFpEF), to study mechanisms of cardiac structural and transcriptional remodeling, cardiomyocyte Ca²⁺ signaling, contraction, and electrophysiology. Outside the lab, he is an avid nature photographer and birdwatcher.



Balazs Horvath, MD, PhD, is an Associate Professor at Department of Physiology, Medical and Health Science Center, University of Debrecen, Debrecen, Hungary. His research focuses on studying molecular and cellular mechanisms of cardiac arrhythmias.



Leighton T. Izu, Ph.D. is Professor at the Department of Pharmacology, University of California, Davis, USA. He studies mathematics, psychology, and biophysics. His research interests include developing mathematical methods for modeling complex systems, understanding how the heart adapts to changing mechanical loads, and studying how variation is beneficial to the population.



John Shaw is a Professor of Aerospace Engineering at the University of Michigan (UM) in Ann Arbor and has been a UM faculty member since 1997. He received a B.S. degree in aerospace engineering from UM in 1985, a M.S. degree in computer science from Virginia Polytechnic and State University in 1989, and a Ph.D. degree in Engineering Mechanics from The University of Texas at Austin in 1997. His research focusses on experimental mechanics, instabilities in materials and structures, thermomechanics of adaptive materials (shape memory alloys and polymers) and structures, and cardiac biomechanics. Past awards include a National Science Foundation CAREER award, Office of Naval Research young investigator award, and the 2013 best paper award in the International Journal of Solids and Structures. He holds eight patents and is a fellow of the American Society of Mechanical Engineers.